

Systematic Review

Adoption of Change: A Systematic Review of the Transtheoretical Model

Caitlin Vela, PT, DPT, CSCS¹; Catherine Ortega, PT, EdD, ATC, OCS^{2*}

¹Baptist Health Link, San Antonio, Texas, USA

²Department of Kinesiology, University of Texas at San Antonio, San Antonio, Texas, USA

*Corresponding author

Catherine Ortega, PT, EdD, ATC, OCS

Associate Professor, Department of Kinesiology, University of Texas at San Antonio, San Antonio, Texas, USA; E-mail: Eli.Ortega1608@gmail.com

Article information

Received: September 9th, 2020; Accepted: September 19th, 2020; Published: September 28th, 2020

Cite this article

Vela C, Ortega C. Adoption of change: A systematic review of the transtheoretical model. *Sport Exerc Med Open J*. 2020; 6(1): 27-38. doi: [10.17140/SEMOJ-6-179](https://doi.org/10.17140/SEMOJ-6-179)

ABSTRACT

Introduction

Sedentary lifestyle is a risk factor for life-altering comorbidities. Though the World Health Organization (WHO) and the Centers for Disease Control (CDC) have provided guidelines, 80% of Americans do not get the recommended physical activity (PA) dose per day. Motivation continues to be an important and elusive factor to effect change.

Purpose

Assess the available evidence regarding the application of the transtheoretical model and stages of change theory (TTM-SOC) in the last 10-years to behavior changes for PA.

Methods

Databases including PSYCInfo, ERIC, CINAHL, EBM, DARE, and OVID, were searched with the following key search terms: “Lifestyle Changes” OR “Lifestyle” OR “Active Living” OR “Lifestyle Changes” OR “Physical Activity” OR “Actigraphy” OR “Exercise” OR “Activity Level” AND “Transtheoretical Model” OR “Stages of Change”. Seventy-nine studies fit the inclusion criteria and were assessed for quality and validity using the PEDrO scale for experimental studies and the specialist unit for review evidence (SURE) for cohort investigations.

Results

Of the five (5) interventional studies included, none used all four components of the TTM-SOC, namely, stages of change, decisional balance, processes of change and self-efficacy. Observational studies were assessed with eleven (11) classified as observational analytical and nineteen (19) as observational descriptive.

Conclusion

None of the investigations assessed the full TTM-SOC. As such, there can be no definitive conclusions with regard to the effectiveness of stage-matched interventions to promote a change from sedentary lifestyle to adoption of PA. There is a need for more rigorous research to test the application of TTM-SOC with both physiologic and quantitative measures for PA.

Keywords

Systematic review; Transtheoretical model; Stage of change; Physical activity.

BACKGROUND

It is recommended that adults perform at least 2.5-hours of moderate-intensity aerobic exercise or 1.25-hours of vigorous-intensity activity or some combination of both types of exercise in a week.¹ However, on average, less than five percent of adults

participate in 30-minutes of physical activity per day.² Current evidence suggests that sedentary lifestyles are associated with comorbidities such as obesity, diabetes mellitus and cardiovascular disease,³ as well as poor health outcomes such as hypertension, hyperlipidemia, stroke, metabolic syndrome, osteoporosis, and certain types of cancers.^{4,5} Though the cause of these comorbidities

ties are multifactorial, a key component to development of these chronic diseases is lack of physical activity. Conversely, increased physical exercise is correlated with a decreased risk for developing these diseases.⁶

Change is needed. Decades of research into human behavior has expanded our understanding of motivation which is key to any behavior change. Attempts to apply the transtheoretical model and stages of change theory (TTM-SOC) to physical activity, have resulted in non-conclusive findings to support or negate this application of the model.^{7,8}

Transtheoretical Model Defined

The TTM-SOC, was created as a compilation of psychological theories to explain behavior change as related to addiction (Prochaska et al). The TTM-SOC initially included the components of stages of change (SOC), decisional balance and processes of change (POC) with self-efficacy (SE) subsequently added. As a motivational theory used to guide interventions for change, the transtheoretical model (TTM) describes the SOC as progressions a person must advance through for the purpose of making an effective behavior change. The SOC consist of precontemplation, contemplation, preparation, action, maintenance and for some behaviors, termination.⁹ The TTM purports that each stage aligns with the constructs of decisional balance-the weighing of pros and cons for behavior change; self-efficacy-confidence to make and maintain the change; and processes of change-both cognitive and behavior.⁹

Previous Research

The TTM has been applied with smoking cessation with physicians and other practitioners adopting stage assessment tools and stage matched interventions.¹⁰ Use of the theory has expanded to explain behavior change with regard to exercise and sedentary lifestyles. In a systematic review, authors concluded that there was cautious support for stage matched exercise interventions.¹¹

In a systematic review by Bulley et al,⁷ (preferred reporting items for systematic reviews and meta-analyses (PRISMA) score=10/27), authors concluded that the accuracy of self-assignment for stages of action or maintenance for physical activity was frequently inconsistent with recommended physical activity guidelines. Furthermore, measurements to assess stages of change for exercise were not found to be valid or standardized, suggesting that more research was needed to investigate the validity of this measure.^{7,12}

Validity limitations of the evidence available for the TTM-SOC prior to 2007 include that stage allocation for exercise was linked with self-reported activity, and the physiological parameters (body composition, physical fitness) used are indirect measurements for changes in stage.⁷ Additionally, though not direct measures of exercise or physical activity, body mass index, weight and girth measurements were used to assess stage progression. With regard to outcome measures used and construct validity, future investigations require validation of instruments with more rigorous

methods.^{7,8} The evidence available regarding TTM-based behavior change interventions applied to exercise as moderate physical activity has been focused upon one component, namely, the stages of change, *versus* all four of the components of the model.⁷ An additional recommendation is that the full model should be assessed fully/holistically.⁸

There is a need to assess whether evidence during the most recent decade has provided more rigorous evaluation and application of the TTM-SOC with and its full components with regard to physical activity.

PURPOSE

The purpose of this review was to determine how the TTM-SOC has been applied for adoption of physical activity in the last 10-years.

METHODS

Search Strategy

The following databases were searched and accessed May 2017 with updates January 2018 and May 2019: PSYCInfo, ERIC, CINAHL, EBM, DARE, OVID. Key search terms for this literature review included: “Lifestyle Changes” OR “Lifestyle” OR “Active Living” OR “Lifestyle Changes” OR “Physical Activity” OR “Actigraphy” OR “Exercise” OR “Activity Level” AND “Transtheoretical Model” OR “Stages of Change.” Inclusion criteria consisted of full text availability, English language, peer reviewed investigations published in the last 10-years, relating to Transtheoretical Model Stages of Change and its application to influence health outcomes *via* physical activity were included in the study. Searching the following additional databases, namely, American Periodicals, ARTbibliographies Modern (ABM), British Periodicals, Digital National Security Archive, ebrary, e-books, GeoRef, PAIS Index, Periodicals Archive Online, ProQuest Dissertations & Theses Global, SciTech Premium Collection, nine (9) articles were found. Of these, all non-peer reviewed findings were omitted.

Inclusion criteria: Investigations that were 1) related to the use of the TTM with physical activity, 2) performed with adults, 3) peer-reviewed, 4) published after 2007, and 5) in the english language (Figure 1).

Initial Review

The search dating 2007-2019 yielded 126 articles. Investigations of reliability and/or validity of model constructs were sequestered from experimental design/randomized controlled trials, though all were organized for further review. Following removal of excluded articles and all subsequent duplicates from overlapping databases, a total of seventy-nine (79) relevant articles were assessed for rigor and inclusion (Figure 1). Articles were organized into two categories: 1) observational (n=61), which included descriptive or analytical studies, and 2) interventional (n=18), consisting of randomized controlled and quasi-experimental studies (Table 1).

Figure 1. Investigations Assessment Flowchart




Table 1. Summary of Interventional Studies

Author/Year (Design)	Purpose	Study Limitations	Population
Findorff et al ¹⁵ (RCT)	To determine whether cross-sectional and longitudinal patterns of exercise adoption reported by older women participating in a randomized trial of an exercise-based fall reduction program would conform to the TTM.	Stage of Change was only variable assessed in control group	Female older adults
Jackson et al ¹⁷ (RCT)	To determine behavior change for physical activity in people with type 2 diabetes in differing stages of change, as a result of delivery of TTM exercise intervention.	Does not measure 3 of 4 constructs of TTM*	Adults with Type 2 Diabetes
Fischer et al ¹⁶ (Quasi-experimental)	To determine effectiveness of certified personal trainer services in promoting exercise adherence in female traditional-aged college students	Stage specific approach not used	Female, traditional-aged college students
Prochaska et al ¹⁸ (Quasi-experimental)	Comparison of initial efficacy of motivational interviewing, online TTM-tailored communications, and brief Health Risk Intervention on 4 health risk factors: inactivity, BMI, stress, smoking	Combined with other theoretical models (Health Risk Intervention, Motivational Interviewing); poor recruitment (25%) and retention (70%) rates; does not measure 4 of 4 constructs of TTM	Adult, university employees
Yang et al ¹⁹ (Quasi-experimental)	Application of TTM to test effects of senior elastic band (SEB) exercises on functional fitness in older adults in contemplation, preparation stages of behavioral change	Does not measure 3 of 4 constructs of TTM; significant differences between groups in lower body flexibility at baseline; convenience sampling	Community older adults (65yrs+), who can stand independently with no AD

Measurement Instruments

Investigators applied the PEDro scale¹³ to interventional investigations. Inclusion or exclusion was not based upon PEDro scores exclusively, rather, investigations scored > 6 were reviewed for validity of methodology and if strong, were included. Investigations with PEDro scores < 5 were excluded.

The Specialist Unit for Review Evidence¹⁴ checklist, specific to cohort or cross-sectional, non-experimental design investigations was chosen to assess the non-interventional investigations (Figure 2) using recommendations regarding critical appraisal checklists.¹⁵

RESULTS

Systematic Reviews

No systematic reviews regarding the TTM-SOC and exercise behavior change were found for the past ten years. This lack of synthesis of recent evidence further validates the need for the current systematic review.

havior change were found for the past ten years. This lack of synthesis of recent evidence further validates the need for the current systematic review.

Interventional Studies

Of the eighteen (18) interventional studies meeting the inclusion criteria, six (6) scored > 6/11 on the PEDro Scale, so judged ‘moderate quality.’ Further in-depth analysis of methodology deemed five (5) to be of moderate to high quality evidence and retained. Of these five (5) investigations, three (3) were quasi-experimental and two (2) were randomized controlled trials.¹⁶⁻²⁰ Of these five (5) articles, two (2) clearly assessed all four constructs of the Transtheoretical Model: self-efficacy, decisional balance, stages of change, and processes of change.^{16,17}

Limitations of the intervention studies included that there was a lack of true stage-matching and there was a failure to account for individuals in precontemplation and contemplation

Figure 2. Cardiff University: SURE Checklist

Questions to Assist with the Critical Appraisal of Cohort Studies

For all questions below, “yes/no/can’t tell”

1. Is the study design clearly stated?
2. Does the study address a clearly focused question? Consider: population; exposure (defined and accurately measured); Comparator/control; outcomes
3. Are the setting, locations, and relevant dates provided? Consider: recruitment period; exposure; follow-up & data collection
4. Were the participants fairly selected? Consider: eligibility criteria; sources & selection of participants; method of follow-up; for matched studies – details of matching criteria and number of exposed or unexposed
5. Are participant characteristics provided? Consider if: sufficient details; a baseline table is included
6. Are the measures of exposures & outcomes appropriate? Consider if the methods of assessment are valid and reliable.
7. Was bias considered? E.g. recall or selection bias
8. Is there a description of how the study size was arrived at?
9. Are the statistical methods well described? Consider: how missing data was handled; were potential sources of bias (confounding factors) controlled for; how loss to follow-up was addressed
 - a. Consider threats to internal validity (all 8, including): _____
10. Is information provided on participant flow? Consider if following provided: flow diagram; numbers of participants at each stage; details of drop-outs; details of missing participant data; follow-up time summarized; numbers of outcome events
11. Are the results well described? Consider if: effect sizes, confidence intervals/standard deviations provided; the conclusions are the same in the abstract and full text
12. Is any sponsorship/conflict of interest reported?
13. Finally...did the authors identify any limitations and, if so, are they captured above?

*Specialist Unit for Review Evidence (SURE)¹⁴

Table 2. Summary of Results (Interventional Studies): Stages of Change

Stage of Change		
Author	Finding	Results
Findorff et al ¹⁵ (RCT)	Individuals with TTM intervention demonstrated increased progression through SOC to become exercise adopters	p value<0.001 CI, SD, effect size not avail
Jackson et al ¹⁷ (RCT)	Individuals receiving exercise consultation interviews demonstrated increased success rates of stage progression versus control group with no consultation	p value 0.007 CI, SD, effect size not avail
Fischer et al ¹⁶ (Quasi-experimental)	Within-group comparison of stage of exercise change from T1 to T2 was significant for the control group.	p value<0.001, α=0.05 CI, SD, effect size not avail
Prochaska et al ¹⁸ (Quasi-experimental)	At 6 months, motivational interviewing and TTM-based intervention groups both demonstrated significantly greater proportions of individuals at criteria (exercising moderately 30 minutes/day for at least 5 days/week) for exercise, as compared to health risk intervention only group.*	p value≤0.01 CI, SD, effect size not avail
Yang et al ¹⁹ (Quasi-experimental)	Participant stages were identified in demographic profiles. Progressions were addressed in article discussion, minimal descriptions.	N/A

*Action phase was singularly assessed in the study Prochaska et al.¹⁸ at follow-up post-intervention. This indicated a greater number of individuals meeting criterion of the action stage versus individuals who did not meet this stage.

Table 3. Summary of Results (Interventional Studies): Decisional Balance

Decisional Balance		
Author	Finding	Results
Findorff et al ¹⁵ (N=272)	No difference among control versus intervention groups	p-value>0.05
Jackson et al ¹⁷ (N=34)	Decisional balance not assessed in this study	N/A
Fischer et al ¹⁶ (N=62)	Significant decrease (negative trend) from T1 to T2 for decisional balance among control group in a measure of which the higher the score, the more beneficial the exercise (pros) is perceived to be.	p-value 0.002, Effect size 0.60, CI, SD not avail
	No significant decrease (no negative trend) among experimental group with personal training services in a measure of which the higher the score, the more beneficial the exercise (pros) is perceived to be.	p-value 0.461 CI, SD not avail
Prochaska et al ¹⁸ (N=1400)	Decisional balance not assessed in this study	N/A
Yang et al ¹⁹ (N=169)	Decisional balance not assessed in this study	N/A

Table 4. Summary of Results (Interventional Studies): Processes of Change

Processes of Change		
Author	Finding	Results
Findorff et al ¹⁵ (N=272)	Increased use of behavioral processes was demonstrated in exercise adopters between T1 and T2	p-value<0.001
	Decreased use of cognitive processes was demonstrated in exercise adopters between T1 and T2	p-value<0.001
	Individuals who progressed in SOC but were not yet in action (exercise readiness) demonstrated decrease in cognitive scores	p-value 0.016
Jackson et al ¹⁷ (N=34)	Processes of change not assessed in this study	N/A
Fischer et al ¹⁶ (N=62)	Within-group comparison of exercise mediators at T1 and T2 demonstrated significant decrease (negative trend) among control group for cognitive processes	p-value 0.006, t[30]=2.963 Effect size 0.53 CI, SD not avail
	Within-group comparison of exercise mediators at T1 and T2 demonstrated significant decrease (negative trend) among control group for behavioral processes	p-value<0.001, t[30]=4.497 Effect size 0.81 CI, SD not avail
	Statistically significant decrease (negative trend) among experimental group with personal training services for behavioral processes	p-value 0.25, Effect size 0.43 CI, SD not avail
	No significant decrease (negative trend) for cognitive processes among experimental group with personal training services	p-value .281, t[30]=1.099 Effect size 0.20 CI, SD not avail
Prochaska et al ¹⁸ (N=1400)	Processes of change not measured in this study	N/A
Yang et al ¹⁹ (N=169)	Processes of change not measured in this study	N/A

Table 5. Summary of Results (Interventional Studies)

Self Efficacy		
Author	Finding	Results
-	Self-efficacy was significant predictor of whether a person would become an exercise adopter long-term	p-value<0.05, CI 1.04-1.14, SD, effect size not avail
	In comparison of T1 to T2 at post-intervention, self-efficacy scores for exercise adopters decreased	p-value 0.43
	In comparison of T1 to T2 at post-intervention, self-efficacy scores for exercise readiness group decreased	p-value 0.29 SD, CI, effect size not avail
Jackson et al ¹⁷ (N=34)	Self-efficacy not assessed in this study	N/A
Fischer et al ¹⁶ (N=62)	Within-group comparison of exercise mediators at T1 and T2 demonstrated significant decrease in scheduling self-efficacy for control group	p-value 0.004, Effect size 0.57, CI, SD not avail
	Within-group comparison of exercise mediators at T1 and T2 demonstrated no significant change for task self-efficacy among control group	p-value 0.863, Effect size 0.03 CI, SD not avail
	Within-group comparison of exercise mediators at T1 and T2 demonstrated no significant change for coping self efficacy among control group	p-value 0.059, Effect size 0.35, CI, SD not avail
	No significant decrease (no negative trend) among experimental group with personal training services for coping self-efficacy, scheduling self efficacy, or task self-efficacy which indicates maintenance or progression in this group.	p-value>0.05, CI, SD not avail
Prochaska et al ¹⁸ (N=1400)	Processes of change not measured in this study	N/A
Yang et al ¹⁹ (N=169)	Processes of change not measured in this study	N/A

Table 6. Summary of Results (Interventional Studies): PA Outcome Measures

Outcome Measure of Physical Activity (Objective)		
Author	Finding	Results
Findorff et al ¹⁵ (N=272)	Walking behavior, as measured by daily exercise logs [primary outcome used for categorization of SOC] demonstrated a significant difference for between-group comparison of individuals in experimental group who became exercise adopters more frequently than the control group.	p-value<0.01
Jackson et al ¹⁷ (N=34)	Among TTM-based intervention, experimental group receiving exercise consultation interviews demonstrated significant change from initial measure in PA levels to 6 weeks	p-value<0.01 CI, SD, effect size not avail
	Among TTM-based intervention, experimental group receiving exercise consultation interviews demonstrated significant change in PA levels after 6 weeks as compared to control group PA levels	p-value<0.01 No additional values avail
Fischer et al ¹⁶ (N=62)	Significant difference for experimental group duration of PA as compared to control group	p-value<0.05 No additional values avail
	Experimental group with personal training services demonstrated significantly more positive pattern of exercise behavior as compared to control group (adherence)	p-value 0.028, Contingency coefficient=0.358 CI, SD, effect size not avail
Prochaska et al ¹⁸ (N=1400)	Primary outcome measure to meet criterion for physical activity was stage of change. Objective PA not clearly tracked in this study.	N/A
Yang et al ¹⁹ (N=169)	Objective PA not tracked in this study.	N/A

Table 7. Summary of Results (Interventional Studies): Other Outcome Measures

Outcome Measure		
Author	Finding	Results
Yang et al ¹⁹ (N=169)	With implementation of senior elastic band program, experimental group participants who engaged in the activity demonstrated significant outperformance at all time points (3- and 6-months) versus control group in: lung capacity, cardiopulmonary fitness, upper and lower body flexibility, upper limb muscle power, and lower limb muscle endurance	p-value<0.05, Effect size 0.2, CI, SD not avail
	All 6 variables (lung capacity, cardiopulmonary fitness, upper and lower body flexibility, upper limb muscle power, and lower limb muscle endurance) were significant for experimental group in comparison of pre-test and 6 months later at post-test	p-value<0.001 CI, SD, effect size not avail
	Among control group with no elastic band exercises, lung capacity changed significantly from pre-test to post-test at 3- and 6-months	p-value 0.34, F=3.61, CI, SD, effect size not avail
	Among control group with no elastic band exercises, cardiopulmonary fitness changed significantly from pre-test to post-test at 3- and 6-months	p-value<0.001, F=14.91, CI, SD, effect size not avail
Health Risk Factors		
Prochaska et al ¹⁸ (N=1400)	At 6 months, TTM and motivational interviewing groups both had significantly greater percentage of participants reaching criterion for effective stress management compared to health risk intervention group	p-value<0.01 CI, SD, effect size not avail
	At 6 months, health risk intervention group demonstrated significantly greater number of health behavior risks than motivational interviewing and TTM groups	p-value<0.05 CI, SD, effect size not avail

who did not have any intention to take action towards physical activity (Table 1). Though these investigations purported to test movement through the stages of change, none of these studies provided stage-matched interventions to assess cause-effect, nor tightly controlled the intervention. Therefore, identified investigations maintained common limitations regarding the TTM-SOC noted in prior decades regarding application to physical activity (Tables 2-7).

Stages of Change

All five (5) interventional studies reported on the stages of change, four (4) of which utilized some variant of a self-report questionnaire to measure the SOC for exercise.^{16-18,20} One (1) remaining study lacked clarity with assessment of the stages, though the exercise SOC appeared to be assigned by the researcher if the participant met a criterion for exercise.¹⁹ Between-stage movement was assessed for exercise SOC among three (3) studies.¹⁶⁻¹⁸ Where regression was noted, one (1) study offered strategies and discussions to overcome the relapse in stages.¹⁶ Data was correlated with

its relation to the current SOC in two (2) studies^{16,18} and in one (1) study, subjects were categorized into their SOC based on a measured amount of physical activity.¹⁸

Decisional Balance

Measured in two (2) of the five (5) included studies, decisional balance was assessed for exercise *via* self-report questionnaire.^{16,17} In both of these studies, there were longitudinal comparisons of decisional balance scores throughout the study,^{16,17} and one (1) of these studies correlated decisional balance with a SOC measurement.¹⁶

Self-Efficacy

Two (2) studies measured SE, which was assessed *via* self-report questionnaires.^{16,17} Longitudinal assessments of SE from baseline to follow-up were included in both of these studies,^{16,17} and in one (1) of these, decisional balance was compared with SOC at post-intervention.¹⁶ This study also used data to determine if SE scores at baseline predicted adoption of exercise by post-intervention.¹⁶

Additionally, three types of SE for task, coping, and scheduling were measured in one (1) study *via* a self-report scale.¹⁷

Processes of Change

The POC were measured by self-report questionnaire and assessed by two (2) of the five (5) studies included in this review.^{16,17} Additionally, included in both studies was a longitudinal comparison of the POC from baseline to follow-up.^{16,17}

Quantitative Measures

Objective quantitative data regarding physical activity was collected by three (3) of the five (5) studies in this review.^{16,19,20} However, two (2) of these included self-report daily exercise logs,^{16,19} and one (1) used functional fitness measures.²⁰

Observational Investigations (Descriptive and Analytical)

Sixty-one (61) observational studies, classified as either descriptive or analytical following analyses with the SURE checklist, did

not yield high-level evidence and therefore, conclusions could not be derived from their results. Despite this level of evidence, only the studies of low rigor (n=31) were omitted completely from the qualitative aspect of this review. The remaining thirty (30) observational studies were of low to moderate quality, including eleven (11) observational analytical designs and nineteen (19) observational descriptive designs.

Supplemental tables describing populations and diagnoses investigated. A descriptive assessment has been included to meet the aims regarding identification of populations and topics investigated in the past ten (10) years regarding the TTM (Supplemental Table 8 and Supplemental Table 9).

DISCUSSION

Similar to systematic reviews performed before 2008, this review has combined a quantitative and qualitative approach without meta-analysis due to variation in reported outcomes. Lack of measured and reported effect sizes results in an inability to quantitatively combine results.^{7,8}

Supplemental Table 8. Summary of Findings (Observational Descriptives of Moderate Quality)

Author/Year (Study)	Population/Age/Ethnicity Investigated	Diagnoses?	SURE Score*
Dumith et al ²²	Southern Brazil	N/A	12/14
Kim ²³	Korean [college students]	N/A	10/14
Rogers et al ²⁴	Adult females	Breast cancer	11/14
Basta et al ²⁵	Adults	HIV-positive	9/14
Dunton et al ²⁶	Middle-aged/Community dwelling adults/United States	N/A	10/14
Rhodes et al ²⁷	Employees/Province of Alberta	N/A	14/14
Sørensen et al ²⁸	Norwegian	N/A	9/14
Cardinal et al ²⁹ (Cohort)	University students/South Korea/United States	N/A	9/14
Cengiz et al ³⁰ (Cross-sectional survey)	University students/Turkey	N/A	10/14
Lutz et al ³¹ (Longitudinal)	Undergraduate students/female/United States	N/A	11/14
Bezyak et al ³² (Survey)	Adults	Mental illness (including schizophrenia, schizoaffective disorder, bipolar disorder, other mood disorders, and other psychotic disorders)	9/14
Jiang et al ³³	American Indian/Alaska Native adults	Pre-diabetes	10/14
Kosma et al ³⁴ (Longitudinal)	Adults	Physical disabilities (including multiple sclerosis, spinal cord injuries)	
Malone et al ³⁵	Adults	Physical disabilities and chronic health conditions (Disabilities or chronic conditions considered to be neuromuscular (such as cerebral palsy, Parkinson's disease, spinal cord injury, stroke, traumatic brain injury), orthopedic (such as amputation, arthritis, low back pain, scoliosis), cardiovascular or pulmonary (such as COPD, dyslipidemia, heart disease, hypertension) and multiple (combination of 2+ disabilities or chronic conditions))	
Yildirim et al ³⁶	Turkish /Women in low or high SES neighborhoods	N/A	11/14
Johnson et al ³⁷	Undergraduate students/ Community volunteers/Adults	N/A	10/14
Colangelo ³⁸	Women	N/A	12/14
Duan et al ³⁹	China/Germany/University Students	N/A	11/14
Kaasalainen et al ⁴⁰	Finnish/Men/Low fitness	N/A	9/14

*Cardiff University Critical Appraisal Checklist: Specialist Unit for Review Evidence (SURE)¹⁴

Supplemental Table 9. Summary of Findings (Observational Analyticals)

Author/Year (Study)	Population/Age/Ethnicity Investigated	Diagnoses?	Cardiff University Critical Appraisal Checklist: SURE Score*
Lorentzen et al ⁴¹	Adults in suburban districts of Oslo		12/14
Hellsten et al ⁴²	Underserved/ minority populations/Females		
Paxton et al ⁴³ (Cross-sectional)	Adult women/Hawaii (United States)	N/A	12/14
Dishman et al ⁴⁴ (Longitudinal; Cohort)	Adults/Hawaii (United States)	N/A	9/14
Lippke et al ⁴⁵	"Adults"		9/14
Stoltz et al ⁴⁶	Adults in weight loss groups and activities		9/14
Plow et al ⁴⁷ (Survey)	Adults	Multiple sclerosis	11/14
Fortier et al ⁴⁸	Adults	Type 2 diabetes	11/14
Dishman et al ⁴⁹	University Students/United States	N/A	9/14
Geller et al ⁵⁰	Adult/Hawaii (United States)	N/A	9/14
Bernard et al ⁵¹	University students/France	N/A	13/14

*Cardiff University Critical Appraisal Checklist: Specialist Unit for Review Evidence (SURE)¹⁴

Of the literature conducted in the last ten-years (2007-2019), five (5) interventional type studies were found to be of adequate methodology and quality for this review. Assessing the results described above, the stages of change is the first construct to be analyzed.

Stages of Change

An accurate and consistent assessment of the stages of change construct is important for a true stage-matched intervention. Two (2) of the five (5) studies assessed SOC *via* means which were unvalidated.^{18,19} Two (2) of the remaining studies utilized SOC questionnaires and reported current research on reliability and validity.^{16,17} Additionally, one study utilized an exercise status questionnaire utilized in 1997.²⁰

Overall, for the SOC construct, quasi-experimental studies and randomized controlled trials, alike have not contributed new data regarding stage changes than had been reported prior to 2007. This appears to be due to failure to appropriately stage-match the interventions based on the TTM. Despite the support to the TTM provided by findings listed above from these studies, conclusions cannot be considered valid because within three (3) of the five (5) studies, the interventions and/or methods of assessment were not appropriate, therefore lacked internal validity.

As has been stated, interventions were not applied as proposed by the TTM-SOC. Only one (1) interventional study demonstrated appropriate, stage-matched measures and yielded moderate quality results and recommendations.¹⁶ Two (2) of the studies claimed to be stage matched, however, their methods and procedures were unclear to delineate that cognitive processes were used in contemplation, and behavioral processes in action and maintenance.^{17,18} One (1) of these investigations inappropriately applied an action-phase intervention to all subjects irrespective of the stage to which they were categorized. Specifically, subjects in pre-

contemplation, a traditionally cognitive-based phase of the model, were given an action-phase intervention.¹⁸ Furthermore, one (1) study lacked appropriate stage-matching altogether, with one third of the participants in the contemplation stage performing the exercise intervention,²⁰ rather than receiving interventions to address the cognitive processes needed to move to action.¹⁹ Results from both studies should be taken cautiously, as integrity to the model is questionable. This lack of internal validity limits the overall validity of findings.

Decisional Balance

Decisional balance (DB) was addressed in one (1) randomized controlled trial (RCT) and one (1) quasi-experimental study of the five (5) interventional studies included in this review.^{16,17} The first, by Findorff et al, reported no significant difference between a group with TTM-based intervention and control group. This finding is inconsistent with the theoretical basis of decisional balance within the TTM.¹⁶

Regarding the cumulative recent evidence on decisional balance, little to no support has been shown to favor the construct as it has been proposed with the TTM. Decisional balance has been omitted in trials based on the TTM regarding health behavior change.

Processes of Change

One (1) RCT and one (1) quasi-experimental study measured the POC.^{16,17} According to results found in the RCT,¹⁶ exercise adopters demonstrated increased use of behavioral processes and decreased use of cognitive processes, consistent with the TTM and displaying positive support for the POC construct. In this same study by Findorff et al, individuals who progressed in the stages but were not yet in action (exercise readiness), demonstrated decreased cognitive scores.¹⁶ This additional finding supports the

POC construct proposed in the TTM.

Overall, there has been mixed findings between the two intervention studies regarding the POC construct. Findorff et al. measured these processes to be in support of the TTM, whereas findings by Fischer and Bryant did not.^{16,17} In a continued trend, this construct also appears to be neglected by researchers when implementing TTM-based interventions for behavior change.

Self-Efficacy

One (1) RCT and one (1) quasi-experimental study in this review measured SE for exercise among their participants.^{15,16} The first of these studies found SE to be a significant predictor of whether a person would become an exercise adopter in the long-term.¹⁶ With a comparison between two time periods, the exercise readiness group demonstrated decreased SE at follow-up.¹⁶ Contrary to the TTM, this study also found SE scores for exercise adopters to decrease when measured post-intervention as compared to their prior scores.¹⁶

Self-efficacy is an additional construct which appears to be inconsistently measured with interventions relating to the transtheoretical model. Based on the literature cited above, recent research has mixed findings which both support and negate the construct within the TTM.

Objective Measures of Physical Activity

Three (3) of the five (5) included interventional studies included subjective assessments of physical activity (PA).¹⁶⁻¹⁸ One (1) additional study appeared to consider the 'action' stage of change as its primary measurement of achieving PA criterion.¹⁹ In one RCT, exercise behavior was exhibited among participants of the experimental group, who became exercise adopters significantly more frequently than the control group.¹⁶ In another RCT, an exercise group receiving exercise consultation interviews as a TTM-based intervention, demonstrated significant change in PA levels from baseline to follow-up at 6-weeks.¹⁸ A significant difference between PA levels was also present when comparing the experimental and control groups, congruent with the TTM.¹⁸

The summative data of recent years on objective PA measures with TTM-based interventions is directed towards subjective measurements of PA. Positive support for the model has also been demonstrated, with these subjective measurements, with the association between increased PA levels or exercise adoption and a TTM-based intervention.^{16,18}

Overarching findings of this review allude to a continued omission of multiple constructs of the TTM with its implementation for exercise behavior change. As was found in the research published by Bulley et al⁷ and Hutchinson et al,⁸ the TTM continues to be inconsistently tested within interventional studies with failure to achieve a true stage-matched intervention. Furthermore, the SOC construct continues to be the singularly emphasized measure among interventions despite the previous recommendations for holistic/full implementation of the model.

A limitation of this review is due to the limited strength of methodology among the studies investigated. Moderate was the highest quality of evidence available. Significant limitations to some of the investigations reviewed included improper stage-matching interventions, self-report with recall bias and unclear description of methodologies regarding stage matched interventions. Though it is difficult to use every construct of the model due to required increase of sample size as variables are added, the lack of assessment of the entire model, is a limitation of recent TTM-SOC literature.

Objective measures of PA have not been cited as a construct of the TTM-SOC. However, based on the model's recent application for health behavior change, specifically for exercise, there must be impetus to require a measure beyond self-report for PA. The intervention investigations relied solely on self-report instruments requiring recollection of exercise, thus introducing recall bias.¹⁶⁻¹⁹ While Findorff et al¹⁶ utilized daily exercise logs, one (1) study measured exercise *via* questionnaires based on recollection of past events, and one (1) study inferred patterns of PA based on answers in the Stages of Change Scale.^{17,18} One investigation by Prochaska et al,¹⁹ was unclear in assessment of frequency and duration of exercise per week, although the authors appeared to consider achievement of the action stage of change for PA (30-minutes of exercise, 5-days per week) to be the equivalent to meeting criterion for that behavior. In a scientific effort to minimize bias, recall bias, and other errors associated with thinking, objective measures such as activity trackers and pedometers are a necessity to increase validity of findings.

CONCLUSION

Evidence added to the TTM-SOC model in the past 10-years does not appear to provide additional guidance for the health practitioner regarding a stage-based approach for PA. It appears that promotion of cognitive process in earlier stages and behavior processes in latter stages, as outlined by the TTM, has neither been negated nor supported.

Holistic assessment of all constructs of the TTM, including SOC, POC, DB, and SE for PA, were limited among the moderate to high quality of evidence, peer-reviewed studies published in the last ten years. As such, there can be no definitive conclusions with regard to the effectiveness of stage-matched interventions to promote a change from sedentary lifestyle to adoption of PA. There is a need for more rigorous research to be performed to test the application of this model for this behavior change and use of physiological and quantitative, measures for PA within future investigations.

Research conducted to-date has assessed behavior change, or adoption of the behavior of PA. Theoretically, the TTM-SOC was designed to explain the cessation of behaviors (Prochaska et al²¹). The application of the TTM-SOC to adoption of PA, appears to bypass cessation and move to adoption. Mechanism must be understood in order to measure and then manage a diagnosis or a process. Therefore, the application of the TTM-SOC to behavior adoption requires more specific investigation, such that the

mechanism of cessation is considered with then the adoption of PA. Only then, would guidance for practitioners be translated with merit from theory to practice.

Therefore, future research is needed with assessment of SOC to cease sedentary lifestyle in addition to the SOC for PA adoption. A true test of the model with stage-matched interventions and use of all constructs must be done for validation with PA.

In an attempt to manage PA lifestyle and PA itself, researchers must devise a method of measurement with high validity and reliability. This should be possible with the current technology available, however, if PA continues as self-assessment, or unassessed, the evidence for the support of the TTM-SOC will remain questionable and of low evidence.

CONFLICTS OF INTEREST

The authors declare that they have no conflicts of interest.

REFERENCES

- Physical Activity Guidelines for Americans. Web site. health.gov. <https://health.gov/paguidelines/second-edition/>. Accessed April 13, 2019.
- Bureau of Labor Statistics. Standing or walking versus sitting on the job in 2016. Web site bls.gov. <https://www.bls.gov/opub/ted/2017/mobile/standing-or-walking-versus-sitting-on-the-job-in-2016.htm>. Accessed April 13, 2019.
- HHS Office and Council on Sports. Facts & Statistics. Web site hhs.gov. <https://www.hhs.gov/fitness/resource-center/facts-and-statistics/index.html>. Accessed April 13, 2019.
- Centers for Disease Control and Prevention (CDC). New CDC report: More than 100 million Americans have diabetes or pre-diabetes. Web site. cdc.gov. <https://www.cdc.gov/media/releases/2017/p0718-diabetes-report.html>. Accessed April 13, 2019.
- U.S. National Library of Medicine. Health Risks of an Inactive Lifestyle. Web site medlineplus.gov. <https://medlineplus.gov/healthrisksofaninactivelifestyle.html>. Accessed April 13, 2019.
- Nigg CR, Geller KS, Motl RW, Horwath CC, Wertin KK, Dishman RK. A research agenda to examine the efficacy and relevance of the transtheoretical model for physical activity behavior. *Psychol Sport Exerc.* 2011; 12(1): 7-12. doi: 10.1016/j.psychsport.2010.04.004
- Bulley C, Donaghy M, Payne A, Mutrie N. A critical review of the validity of measuring stages of change in relation to exercise and moderate physical activity. *Critical Public Health.* 2007; 17(1): 17-30. doi: 10.1080/09581590601045261
- Hutchison AJ, Breckon JD, Johnston LH. Physical activity behavior change interventions based on the transtheoretical model: A systematic review. *Health Educ Behav.* 2009; 36(5), 829-845. doi: 10.1177/1090198108318491
- Prochaska JO, Redding CA, Evers KE. The transtheoretical model and stages of change. In: Health Behavior and Health Education: Theory, Research, and Practice. 4th ed. San Francisco, CA, USA: Jossey-Bass; 2008: 97-121.
- U.S. Public Health Service. A clinical practice guideline for treating tobacco use and dependence. *J Am Med Assoc.* 2000; 283: 3244-3254.
- Spencer L, Adams TB, Malone S, Roy L, Yost E. Applying the transtheoretical model to exercise: A systematic and comprehensive review of the literature. *Health Promotion Practice.* 2006; 7(4): 428-443. doi: 10.1177/1524839905278900
- Marshall SJ, Biddle SJ. The transtheoretical model of behavior change: A meta-analysis of applications to physical activity and exercise. *Ann Behav Med.* 2001; 23(4): 229-246. doi: 10.1207/S15324796ABM2304_2
- PEDro scale. Web site pedro.org. <https://www.pedro.org.au/english/downloads/pedro-scale/>. Accessed April 13, 2019.
- Specialist Unit for Review Evidence (SURE). Questions to assist with the critical appraisal of cohort studies. Web site. <http://www.cardiff.ac.uk/insrv/libraries/sure/checklists.html>. Accessed April 13, 2019.
- Sanderson S, Tatt ID, Higgins J. Tools for assessing quality and susceptibility to bias in observational studies in epidemiology: A systematic review and annotated bibliography. *Int J Epidemiol.* 2007; 36(3): 666-676. doi: 10.1093/ije/dym018
- Findorff MJ, Hatch Stock H, Gross CR, Wyman JF. Does the transtheoretical model (TTM) explain exercise behavior in a community-based sample of older women? *J Aging Health.* 2007; 19(6): 985-1003. doi: 10.1177/0898264307308341
- Fischer DV, Bryant J. Effect of certified personal trainer services on stage of exercise behavior and exercise mediators in female college students. *J Am Coll Health.* 2008; 56(4): 369-376. doi: 10.3200/JACH.56.4.369-376
- Jackson R, Asimakopoulou K, Scammell A. Assessment of the transtheoretical model as used by dietitians in promoting physical activity in people with type 2 diabetes. *J Hum Nutr Diet.* 2007; 20(1): 27-36. doi: 10.1111/j.1365-277X.2007.00746.x
- Prochaska JO, Butterworth S, Redding CA, Burden V, Perrin N, Leo M, et al. Initial efficacy of MI, TTM tailoring and HPIs with multiple behaviors for employee health promotion. *Prev Med.* 2008; 46(3): 226-231. doi: 10.1016/j.ypmed.2007.11.007
- Yang HJ, Chen KM, Chen MD, Wu HC, Chang WJ, Wang YC, et al. Applying the transtheoretical model to promote functional fitness of community older adults participating in elastic band exer-

- cises. *J Adv Nurs*. 2015; 71(10): 2338-2349. doi: 10.1111/jan.12705
21. Prochaska JO, DiClemente CC. Stages and processes of self-change of smoking: Toward an integrative model of change. *J Consult Clin Psychol*. 1983; 51(3): 390-395. doi: 10.1037//0022-006x.51.3.390
22. Dumith SC, Gigante DP, Domingues MR. Stages of change for physical activity in adults from Southern Brazil: A population-based survey. *Int J Behav Nutr Phys Act*. 2007; 4(1): 25. doi: 10.1186/1479-5868-4-25
23. Kim YH. Application of the transtheoretical model to identify psychological constructs influencing exercise behavior: A questionnaire survey. *Int J Nurs Stud*. 2007; 44(6): 936-944. doi: 10.1016/j.ijnurstu.2006.03.008
24. Basta TB, Reece M, Wilson MG. The transtheoretical model and exercise among individuals living with HIV. *Am J Health Behav*. 2008; 32(4): 356-367. doi: 10.5555/ajhb.2008.32.4.356
25. Rogers LQ, Courneya KS, Shah P, Dunnington G, Hopkins-Price P. Exercise stage of change, barriers, expectations, values and preferences among breast cancer patients during treatment: A pilot study. *Eur J Cancer Care (Engl)*. 2007; 16(1): 55-66. doi: 10.1111/j.1365-2354.2006.00705.x
26. Dunton GF, Vaughan E. Anticipated affective consequences of physical activity adoption and maintenance. *Health Psychol*. 2008; 27(6): 703-710. doi: 10.1037/0278-6133.27.6.703
27. Rhodes RE, Plotnikoff RC, Courneya KS. Predicting the physical activity intention-behavior profiles of adopters and maintainers using three social cognition models. *Ann Behav Med*. 2008; 36(3): 244-252. doi: 10.1007/s12160-008-9071-6
28. Sørensen M, Gill DL. Perceived barriers to physical activity across Norwegian adult age groups, gender and stages of change. *Scand J Med Sci Sports*. 2008; 18(5): 651-663. doi: 10.1111/j.1600-0838.2007.00686.x
29. Cardinal BJ, Lee JY, Kim YH, Lee H, Li KK, Si Q. Behavioral, demographic, psychosocial, and sociocultural concomitants of stage of change for physical activity behavior in a mixed-culture sample. *Am J Health Promot*. 2009; 23(4): 274-278. doi: 10.4278/ajhp.06051667
30. Cengiz C, Çiçek ML. Exercise stages of change in Turkish university students by sex, residence, and department. *Perceptual and Motor Skills*. 2009; 108(2): 411-421. doi: 10.2466/PMS.108.2.411-421
31. Lutz RS, Stults-Kolehmainen MA, Bartholomew JB. Exercise caution when stressed: Stages of change and the stress-exercise participation relationship. *Psychology of Sport and Exercise*. 2010; 11(6): 560-567. doi: 10.1016/j.psychsport.2010.06.005
32. Bezyak JL, Berven NL, Chan F. Stages of change and physical activity among individuals with severe mental illness. *Rehabil Psychol*. 2011; 56(3): 182-190. doi: 10.1037/a0024207
33. Jiang L, Beals J, Zhang L, Mitchell CM, Manson SM, Acton KJ, et al. Latent class analysis of stages of change for multiple health behaviors: Results from the special diabetes program for indians diabetes prevention program. *Prev Sci*. 2012; 13(5): 449-461. doi: 10.1007/s11121-011-0272-z
34. Kosma M, Ellis R, Bauer JJ. Longitudinal changes in psychosocial constructs and physical activity among adults with physical disabilities. *Disability and Health Journal*. 2012; 5(1): 1-8. doi: 10.1016/j.dhjo.2011.09.002
35. Malone LA, Barfield J, Brasher JD. Perceived benefits and barriers to exercise among persons with physical disabilities or chronic health conditions within action or maintenance stages of exercise. *Disabil Health J*. 2012; 5(4): 254-260. doi: 10.1016/j.dhjo.2012.05.004
36. Yildirim, G, Ince ML, Muftuler M. Physical activity and perceptions of Neighborhood walkability among Turkish women in low and high socio-economic Environments: An exploratory study. *Percept Mot Skills*. 2012; 115(2): 661-675. doi: 10.2466/06.15.PMS.115.5.661-675
37. Johnson P, Fallon EA, Harris BS, Burton B. Body satisfaction is associated with Transtheoretical model constructs for physical activity behavior change. *Body Image*. 2013; 10(2): 163-174. doi: 10.1016/j.bodyim.2012.12.002
38. Colangelo AM. *Supportive Exercise: Exploring the Relationship of Social Support and Stages of Change in Relationship to Exercise Behavior in Women*. [dissertation]. Washington, DC, USA; American University; 2014.
39. Duan Y, Brehm W, Wagner P, Chung PK, Graf S, Zhang R, et al. Transition to adulthood: Relationships among psychosocial correlates, stages of change for physical activity, and health outcomes in a cross-cultural sample. *J Phys Act Health*. 2015; 12(11): 1461-1468. doi: 10.1123/jpah.2014-0389
40. Kaasalainen KS, Kasila K, Komulainen J, Malvela M, Poskiparta M. Readiness for health behavior changes among low fitness men in a finnish health promotion campaign. *Health Promot Int*. 2015; 31(4): 827-838. doi: 10.1093/heapro/dav068
41. Lorentzen C, Ommundsen Y, Holme I. Psychosocial correlates of stages of change in physical activity in an adult community sample. *European Journal of Sport Science*. 2007; 7(2): 93-106. doi: 10.1080/17461390701456122
42. Hellsten LA, Nigg C, Norman G, Burbank P, Braun L, Breger R, et al. Accumulation of behavioral validation evidence for physical activity stage of change. *Health Psychol*. 2008; 27(1S): S43-S53. doi: 10.1037/0278-6133.27.1(Suppl.).S43
43. Paxton RJ, Nigg CR, Motl RW, McGee K, McCurdy D, Matthai,

- CH, et al. Are constructs of the transtheoretical model for physical activity measured equivalently between sexes, age groups, and ethnicities?. *Ann Behav Med.* 2008; 35(3): 308-318. doi: [10.1007/s12160-008-9035-x](https://doi.org/10.1007/s12160-008-9035-x)
44. Dishman RK, Thom NJ, Rooks CR, Motl RW, Horwath C, Nigg CR. Failure of post-action stages of the transtheoretical model to predict change in regular physical activity: A multiethnic cohort study. *Ann Behav Med.* 2009; 37(3): 280-293. doi: [10.1007/s12160-009-9113-8](https://doi.org/10.1007/s12160-009-9113-8)
45. Lippke S, Plotnikoff RC. The protection motivation theory within the stages of the transtheoretical model—Stage-specific interplay of variables and prediction of exercise stage transitions. *Br J Health Psychol.* 2009; 14(2): 211-229. doi: [10.1348/135910708X399906](https://doi.org/10.1348/135910708X399906)
46. Stoltz KB, Reysen RH, Wolff LA, Kern RM. Lifestyle and the stages of change in weight loss. *Journal of Individual Psychology.* 2009; 65(1): 69-85.
47. Plow MA, Finlayson M, Cho C. Correlates of stages of change for physical activity in adults with multiple sclerosis. *Res Nurs Health.* 2011; 34(5): 378-388. doi: [10.1002/nur.20447](https://doi.org/10.1002/nur.20447)
48. Fortier MS, Sweet SN, Tulloch H, Blanchard CM, Sigal RJ, Kenny GP, et al. Self-determination and exercise stages of change: Results from the diabetes Aerobic and resistance exercise trial. *J Health Psychol.* 2012; 17(1): 87-99. doi: [10.1177/1359105311408948](https://doi.org/10.1177/1359105311408948)
49. Dishman RK, Jackson AS, Bray MS. Validity of processes of change in physical activity among college students in the TIGER study. *Ann Behav Med.* 2010; 40(2): 164-175. doi: [10.1007/s12160-010-9208-2](https://doi.org/10.1007/s12160-010-9208-2)
50. Geller KS, Nigg CR, Motl RW, Horwath CC, Dishman RK. Transtheoretical model constructs for physical activity behavior are invariant across time among ethnically diverse adults in Hawaii. *Psychol Sport Exerc.* 2012; 13(5): 606-613. doi: [10.1007/s12160-010-9208-2](https://doi.org/10.1007/s12160-010-9208-2)
51. Bernard P, Romain AJ, Trouillet R, Gernigon C, Nigg C, Ninot G. Validation of the TTM processes of change measure for physical activity in an adult French sample. *Int J Behav Med.* 2014; 21(2): 402-410. doi: [10.1007/s12529-013-9292-3](https://doi.org/10.1007/s12529-013-9292-3)