

Case Report

Surgical Treatment of Sternal Fracture: Case Report and Literature Review

Jorge Pérez-Acosta, MD, MSc^{1,2}; Orlando Téllez-Almenares, MD, MSc^{1,2*}; Justo Escalona-Cartaya, MD^{1,2}; Raúl Calas-Balbuena, MD, MSc^{1,2}; Olga M. Moreira-Barinaga, MD¹

¹Saturnino Lora Provincial Hospital, Santiago de Cuba, Cuba

²University of Medical Sciences of Santiago de Cuba, Santiago de Cuba, Cuba

*Corresponding author

Orlando Téllez-Almenares, MD, MSc

Saturnino Lora Provincial Hospital, Santiago de Cuba, Cuba; University of Medical Sciences of Santiago de Cuba, Santiago de Cuba, Cuba;

E-mail: orlandotellez.al@gmail.com

Article information

Received: June 8th, 2023; Revised: July 10th, 2023; Accepted: July 14th, 2023; Published: July 21st, 2023

Cite this article

Pérez-Acosta J, Téllez-Almenares O, Escalona-Cartaya J, Calas-Balbuena R, Moreira-Barinaga OM. Surgical treatment of sternal fracture: Case report and literature review. *Surg Res Open J.* 2023; 8(1): 1-6. doi: [10.17140/SROJ-8-129](https://doi.org/10.17140/SROJ-8-129)

ABSTRACT

Sternal fractures are uncommon thoracic trauma that can ensue in isolation or conjunction with other organ injuries, implying more significant morbidity and mortality. These fractures result from diverse aetiological mechanisms, some of which may provoke minor chest wall issues, while others can have deadly outcomes. The sternal fracture's primary aetiological culprits are direct impact and the upper thoracic region's compression. Studies reveal that traumatic sternum fractures occur in 8-18% of blunt thoracic and polytrauma victims, whereas they are unusual in open trauma. This report aims to outline the case of a 54-year-old Caucasian male with hypertension and an alcohol consumption history who sustained a direct impact on the anterior thorax after a traffic accident. He experienced severe pain that worsened with movement and coughing, and breathing difficulty. We noticed a painful haematoma and depression with bony crepitus at the sternal body level. Abdominal-pelvic echography indicated no intra-abdominal lesions and showed pleural effusion in the left lung base, while thoracic X-rays (lateral and anteroposterior view) and chest computerized tomography (CT) scan exhibited mid-sternum segment displaced fracture and haemothorax. We diagnosed a mid-sternum completely displaced overlapping fracture and a traumatic left haemothorax. Therefore, urgent surgical intervention was required. We performed a minimal-lower pleurotomy and internal fixation to address the fracture. His post-operative recovery was satisfactory. Although various authors advocate X-rays as an effective diagnostic means for sternal fractures, CT uncovers overlooked radiography abnormalities; therefore, CT must be considered in patients' initial assessment. While most sternal fractures can be treated conservatively, surgery is feasible in displaced fractures.

Keywords

Case report; Osteosynthesis; Sternal fracture; Sternum; Thoracic trauma.

INTRODUCTION

Trauma remains a global health concern with meaningful repercussions on peoples' lives and healthcare systems.^{1,2} The development and mechanisation of industries and vehicles' widespread use has been determining factors in increased traumas.³ The Swedish Trauma Registry informs that it is one of the leading cause of death and disability in individuals under 45 worldwide, surpassing cancer deaths among young adults.⁴

Road traffic crashes are a foremost cause of injury, posing a critical matter in numerous nations.² The World Health Organisation (WHO) conveyed that traffic accidents provoke approximately 1.3 million yearly casualties.⁵ As per National Trauma Data Bank's 2016 annual report,⁶ 223,866 motor vehicle accidents ensued in the

United States, resulting in 10,343 casualties. A recent Cuban report⁷ points out that between 2019 and 2020, there were 1,368 fatalities from road traffic accidents, of which 80.8% were male.

Blunt trauma is a common occurrence in motor vehicle collisions and is associated, in most cases, with severe organ damage.^{1,6,8} In Sweden, blunt injuries account for approximately 90% of all injuries. Blunt thorax traumas (BTT) are the third leading cause of trauma-related death, preceded only by traffic-related head and abdominal injuries.⁸

The thorax is one of the most affected regions in motor vehicle accidents. The 2016 National Trauma Data Bank statistics⁶ logged 1,29,338 thoracic traumas (TT) with AIS>3 with a fatality rate of 9.53. Multiple authors have noted that TT mortality is ap-

proximately 25% of all traumatic deaths globally and contributes 25% to mortality from other types of traumas. South Korean research Byun et al¹ shows that 33.6%-non-surviving caseload died due to significant chest harm. In contrast, a Cuban study³ states that TT accounts for 4-6% of trauma admissions in Cuba and has a fatality rate of 15%.

The sternum is a peculiar, odd bone located in the anteromedial thorax region, it measures 15-20 cm in length, and as it extends distally, its thickness gradually decreases.⁹ Fractures of this bone are uncommon and can ensue in isolation or alongside other organ damage, implying more significant morbidity and mortality.^{10,11} Studies Brookes et al,¹² Hochhegger et al,¹³ Knobloch et al,¹⁴ and Bentley et al¹⁵ reveal traumatic sternal fractures (SFs) occur in 8-18% of BTT and polytrauma victims. In contrast, they are unusual in open trauma and refer that using seat belts is associated with a higher frequency of these fractures.

These fractures arise from diverse aetiological mechanisms, some of which may provoke minor chest wall injuries, while others can have deadly outcomes. The sternal fracture's main aetiological culprits are direct impact, the upper thoracic region's compression, and deceleration.^{16,17} In car crashes, the SFs are generally associate with seat belt use or the blow against the steering wheels.¹⁷

This report aims to outline the case of a patient who suffered a traumatic SF after a traffic crash, the injury's characteristics, and its clinical and surgical management.

CASE PRESENTATION

Clinical Assessment

A 54-year-old Caucasian male with hypertension and an alcohol consumption history was transferred by the Integrated Medical Emergency Service (SIUM for its Spanish acronym) to the Accident and Emergency Department (A&E) following a traffic incident. He sustained a direct impact on the anterior thorax, causing severe pain that worsened with movement and coughing, making it difficult for him to breathe.

In the medical interview, the patient shared accident details describing how he collided with a parked lorry while riding his motorbike. He confirmed the impact was only in his chest and denied experiencing discomfort in his abdomen, neck or other regions.

We adhered to the advanced trauma life support (ATLS) guidelines, and a complete physical examination was conducted. Two peripheral veins were cannulated, and a Fowley catheter was placed.

The noteworthy findings on the physical assessment included a weight of 65 kg, a height of 172 cm, and a body mass index (BMI) of 22.4 kg/m²; the skin and mucous membranes appeared slightly pale and dry, predominantly affecting the conjunctival and oral mucosa. We noticed a haematoma and depression at the sternal body level. It was painful on palpation and had bony

crepitus. No abnormalities were detected upon abdominal examination through palpation, percussion, and auscultation. Additionally, the abdominal puncture was negative.

Upon respiratory system asses, it was found that there was reduced thoracic expansion, a respiratory rate of 28 breaths per minute, decreased vocal resonance, and vesicular breath sounds towards the left lung base without crackles or pleural friction rub. As for the cardiovascular system, was found audible heart sounds, good tone with tachycardic, and a heart rate (HR) of 114 beats per min (b.p.m). Peripheral pulses were present, accelerated and synchronous, and blood pressure was measured at 140/110 mmHg.

Neurological sphere assessment showed that consciousness and orientation were preserved, and a Glasgow Scale score of 15 points was logged.

During the patient's initial evaluation at A&E, we diagnosed mild dehydration. We were administered crystalloid fluids (isotonic sodium chloride) at a 1,500 ml/m²sc/24 h rate, passing 750 ml of fluid in the first hour. We conducted paraclinical examinations after ensuring the patient's clinical and haemodynamic stability.

Laboratory Tests

The detailed values of all the laboratory tests are given in the below box.

Box. Laboratory Tests Values
Partial blood red cell measurements: Hb (135 g/L (13.5 g/dL), Ht. (40.5 L/L)
Erythrocyte sedimentation rate (ESR) (5 mm/h)
Leukogram:
leukocytes (7x10 ⁹ /L)
neutrophils: 0.65
eosinophils: 0.01
lymphocytes: 0.31
monocytes: 0.03
basophils: 0.00
Partial screening test of haemostasis:
Clotting time 7 min
Bleeding time 2 min
Platelet count (270x10 ⁹ /L)
Retractile clot. Glycaemia (5.8 mmol/L)
Arterial blood gasses analysis:
pH (7.40)
PaCO ₂ (38 mmHg)
PaO ₂ (97 mmHg)
HCO ₃ (26 mEq)
K ⁺ (4.0 mmol/L)
Na ⁺ (143 mmol/L)
Cl ⁻ (98 mmol/L)

Electrocardiogram

Electrocardiogram showed sinus tachycardia without changes in the Q wave, R wave and S wave (QRS complex) or ST-segment and HR of 112 b.p.m.

Imaging Studies

Abdominal-pelvic echography showed no organ damage, no fluid in the abdominal cavity, no urinary tract injury, and an empty bladder due to the Fowley catheter inside. Nevertheless, there was evidence of pleural effusion in the left lung base.

Chest X-ray in a standing position (posteroanterior view) showed sternum body displaced fracture and a radiopacity that obscured the left costophrenic angle. (Figures 1 and 2).

Figure 1. Sagittal CT Scan Image: Mid-sternum Completely Displaced Overlapping Fracture



Figure 2. Coronal CT Scan Images: The Mid-sternum Completely Displaced Overlapping Fracture




Surgical Management and Post-surgery

Based on clinical and imaging studies, it was determined that the patient had suffered a mid-sternum complete displaced overlapping fracture and a traumatic left haemothorax. As a result, emergency surgical treatment was required.

The haemothorax was evacuated through a minimal lower pleurotomy under local anaesthesia; it removed approximately 480 ml of haematic contents. The sternum was fixed under general endotracheal anaesthesia using an internal fixation system of galvanised stainless-steel plates and screws (Figure 3).

Figure 3. A. The Incision Made Over the Sternum and the Sternal Fracture is shown B. Placement of the Fixation System for Osteosynthesis




The fracture was accessed through a longitudinal incision approximately 12-15 cm above the sternum. Fortunately, there were no intraoperative complications.

Following a 1 hour and 20-minute surgery, he was transferred to the intermediate care unit (ICU), where he stayed for 12-days. During his stay, he experienced cardiac arrhythmia, which subsided after removing his chest tube. After 18-days, he was discharged with regular outpatient follow-up and pain management.

The patient complained of mild pain three months later, but the sternum was stable. There were no reported complications at six months, and the radiological images showed favourable healing with intact bony fixation (Figure 4).

Figure 4. Post-operative Evolutionary Chest X-rays: A. Trans-operative; B and C. Evolutionary in ICU; D, E and F. Evolutionary in Consultation



DISCUSSION

Most of the reviewed-available literature indicates sternum fractures are infrequent TT. However, authors such as Bentley et al¹⁵ argue that it is not as uncommon as believed. Some researchers^{18,19} claim that SFs diagnoses have increased recently because of the increasingly routine use of computerized tomography (CT) scans in multiple trauma patients' initial assessment. Şimşek et al²⁰ con-

tend that SFs account for a small thoracic trauma proportion and occur in 4% of road traffic accidents and 3-8% of blunt abdominal traumas. Meanwhile, Doyle et al¹⁹ and Hochhegger et al¹³ based on previous research, inform 68% of SFs incidence in motor vehicle collisions, 7.9% in falls and motorbike accidents, respectively, 3.4% in pedestrian-struck, and 1.4% in bicycle accidents.

Ayes-Valladares et al²¹ apprise that SFs secondary to blunt chest trauma are rare in Honduras, where only a 1.4% incidence rate is notified. Similarly, reports Fonseca Sosa,¹⁰ Ferrer Lozano et al,¹⁶ Nazario Dolz et al²² and Herrera et al²³ indicate that Cuba's frequency rate is low; i.e., Moya Rosa et al²⁴ reported a rate of 0.78% in their series. Isolated cases of SFs have been reported in our country. One such case involved a subject gored by an ox, as Nazario et al²² documented from our hospital.

Sternal fractures are typically caused by direct impacts or deceleration, resulting in sternal body or manubrium harm, by and large. Fracture severity varies depending on the accompanying lesions.^{15,16} Commonly, fractures occur at the sternal body level and in its distal portion due to the tapering of the bone in the caudal direction. According to the Brookes et al¹² caseload and von Garrel et al²⁵ study most lesions (93.4% and 76.5%, respectively) were found in the sternal body. Notwithstanding, Şimşek's et al²⁶ research of 108 subjects revealed that 59.3% had injuries in the sternal manubrium, while only 39.1% had them in the sternal body.

The SFs diagnosis is primarily based on the history recorded by interrogation, clinical findings on physical examination and imaging studies. Suspicion signs comprise a trauma history, retrosternal pain exacerbated by coughing or inspiration, respiratory distress, oedema, ecchymosis or haematomas at the sternum level. Other findings include bone crepitus and depression.^{13,15,26}

Detecting STs through anteroposterior radiography alone has a sensitivity of only 50%; nevertheless, when combined with a lateral view chest X-ray, sensitivity increases because these fractures typically occur in the sagittal plane.^{15,24,27}

Some authors, Ferrer Lozano et al¹⁶ and Nazario Dolz et al²² especially in our environment, state that the lateral and anteroposterior chest X-ray views provide an adequate STs initial diagnosis, whereas a CT scan should only be used if additional radiological abnormalities are detected. Nonetheless, while radiography can still be a valid diagnostic option, other experts Perez et al,¹⁸ Doyle et al,¹⁹ Ramgopal et al²⁷ and Kim et al²⁸ assert that CT scans are the gold standard for detecting sternal fractures and should be required. We align with this standpoint.

It is crucial excluding associated injuries in these patients. It is rare for isolated sternal fractures to be related to blunt cardiac traumas. Regardless, if they do occur, the mortality rate is 0.8%. In polytrauma cases, concurrent injuries can be severe and lead to a mortality rate as high as 7.9%.^{29,30}

In Şimşek's et al²⁶ the most frequent intrathoracic lesions were pulmonary contusion and pneumothorax (53.7% each), while pneumothorax occurred in 51.9%. In contrast, in Perez's report,¹⁸

the highest intrathoracic injury incidence was a pulmonary contusion (26.3%), followed by pneumothorax (23.6%), while haemothorax had an incidence of 11.9%.

Acute sternal trauma management should adhere to the guidelines set out in the ATLS.¹⁵ Identifying associated life-threatening injuries should be a priority to rule out damages such as cardiac tamponade (uncommon in closed chest trauma), tension pneumothorax, large haemothorax or other significant damages to the skull, abdomen and pelvis.

SFs surgical treatment is infrequent, and its initial choice is controversial; conservative management is generally chosen.^{11,18,31,32} Regardless, surgical treatment should be used in intractable pain cases, altered ventilatory mechanics, and poor bony alignment to optimise rehabilitation due to rachimedullary trauma.^{18,33}

McKim made the first sternal fracture fixation description. Over time, various materials for fixation have appeared, but the most widely accepted technique is plate osteosynthesis using titanium plates and screws.¹⁹ However, Harston et al,³⁴ in his meta-analysis, where he analysed 52 individuals who underwent fixation with plates and 24 with wire, showed healing in all cases regardless of the technique used.

CONCLUSION

Although various authors advocate X-rays as an effective diagnostic means for sternal fractures, CT uncovers overlooked radiography abnormalities; therefore, CT must be considered in patients' initial assessment. While most sternal fractures can be treated conservatively, surgery is feasible in displaced fractures.

PATIENT PERSPECTIVE

The patient confirms that he is pleased with the treatment and aftercare provided by the medical staff. He acknowledges that he was well-informed about his injury, its potential complications, and the surgical procedure he underwent. He feels content with his progress and has gradually resumed his daily routines.

ETHICAL RESPONSIBILITIES

Protection of People and Animals

The authors declare that no experimentation was performed on the patient, and therapeutic management was under our hospital standards.

Data Confidentiality

No data that could compromise the patient's privacy is disclosed in this report.

RESEARCH QUALITY

The authors adhered to the EQUATOR Guidelines recommen-

dations and specifically to CARE Guidelines for publishing Case Reports.

FINANCIAL SUPPORT AND SPONSORSHIP

The authors stated that no grants were involved in supporting this work.

ACKNOWLEDGEMENTS

We thank the radiologists and intensivists, the nurses and the operating theatre technical staff involved in the patient's care.

CONSENT

The patient consented to use his medical history data in this report.

CONFLICTS OF INTEREST

No competing interests were disclosed.

REFERENCES

- Byun CS, Park IH, Oh JH, Bae KS, Lee KH, Lee E. Epidemiology of trauma patients and analysis of 268 mortality cases: Trends of a single Center in Korea. *Yonsei Med J.* 2015; 56(1): 220-226. doi: 10.3349/ymj.2015.56.1.220
- Beshay M, Mertzluft F, Kottkamp HW, et al. Analysis of risk factors in thoracic trauma patients with a comparison of a modern trauma centre: a mono-centre study. *World J Emerg Surg.* 2020; 15: 45. doi: 10.1186/s13017-020-00324-1
- Satorre Rocha JA, León González OC, López Rodríguez PR, García Castillo E, Danta Fundora LM. Characterisation of patients with thoracic trauma treated at the Hospital Enrique Cabrera. 2014-2018. *MediSur.* 2019; 17(6): 815-823.
- SweTrau. Svenska Traumaregistret. Website. <https://rcsyd.se/swetrau/>. Accessed July 7, 2023.
- World Health Organization (WHO). Road traffic injuries. Website. <https://www.who.int/news-room/fact-sheets/detail/road-traffic-injuries>. Accessed July 3, 2023.
- American College of Surgeons (ACS). NTDB Reports and Publications. Website. <https://www.facs.org/quality-programs/trauma/quality/national-trauma-data-bank/reports-and-publications/>. Accessed July 7, 2023.
- Ministry of Public Health. Health Statistical Yearbook 2020. Website. <https://temas.sld.cu/estadisticassalud/>. Accessed March 6, 2023.
- Lundin A, Akram SK, Berg L, Göransson KE, Enocson A. Thoracic injuries in trauma patients: Epidemiology and its influence on mortality. *Scand J Trauma Resusc Emerg Med.* 2022; 30(1): 69. doi: 10.1186/s13049-022-01058-6
- Latarjet M, Ruíz Liard A, Pró E. Sección XIII. Capítulo 71. Caja Torácica. In: *Anatomía Humana*. [In: Spanish]. 5^{ta} ed. España, Spain: Editorial Médica Panamericana; 2019: 860.
- Fonseca Sosa FK. Sternal fracture. Presentation of a case. *Rev Chil Cir.* 2017; 69(5): 408-411. doi: 10.1016/j.rchic.2016.10.018
- Tinajero JCJ, Santillán-Doherty P, Argote-Greene LM, et al. Traumatic fractures of the sternum: Priorities and management strategy. *NCT Neumol Cir Torax.* 2009; 68(4): 151-155.
- Brookes JG, Dunn RJ, Rogers IR. Sternal fractures: A retrospective analysis of 272 cases. *J Trauma.* 1993; 35(1): 46-54. doi: 10.1097/00005373-199307000-00008
- Hochegger B, Altmayer S. Traumatic sternal fractures. *Radiol Bras.* 2022; 55(4): IX. doi: 10.1590/0100-3984.2022.55.4e3
- Knobloch K, Wagner S, Haasper C, et al. Sternal fractures occur most often in old cars to seat-belted drivers without any airbag often with concomitant spinal injuries: Clinical findings and technical collision variables among 42,055 crash victims. *Ann Thorac Surg.* 2006; 82(2): 444-450. doi: 10.1016/j.athoracsur.2006.03.046
- Bentley TP, Ponnarasu S, Journey JD. Sternal Fracture. In: *StatPearls*. Treasure Island (FL), USA: StatPearls Publishing; 2022.
- Ferrer Lozano Y, Oquendo Vazquez P, Morejón Trofimova Y. Isolated sternal fracture. Presentation of a case. *MediSur.* 2014; 12(6): 889-894.
- Klei DS, Schutte H, Öner FC, van Baal MC, Leenen LP, van Wessem KJ. Traumatic sternal fractures can be safely treated conservatively - A 13-year retrospective cohort study. *J Surg Res (Houst).* 2021; 4(4): 572-587. doi: 10.26502/jsr.10020170
- Perez MR, Rodriguez RM, Baumann BM, et al. Sternal fracture in the age of pan-scan. *Injury.* 2015; 46(7): 1324-1327. doi: 10.1016/j.injury.2015.03.015
- Doyle JE, Diaz-Gutierrez I. Traumatic sternal fractures: A narrative review. *Mediastinum.* 2023; 5: 34. doi: 10.21037/med-21-27
- Şimşek S, Özmen CA, Onat S. Morbidity and mortality associated with fracture of the sternum due to blunt trauma, by fracture type and location. *Radiol Bras.* 2022; 55(3): 167-172. doi: 10.1590/0100-3984.2021.0074
- Ayes-Valladares F, Alvarado LT. Clinical and therapeutic characterisation of sternal fractures in the Hospital Escuela. *Rev Méd Hondur.* 2009; 77(3): 114-117.
- Nazario Dolz AM, Orosco García L, Domínguez González EJ, Rodríguez Fernández Z, Falcón Vilariño CG. Sternal fracture due to ox goring. *Rev Cuba Cir.* 2013; 52(4): 315-323.

23. Herrera JM, Soler J de JM. Sternal fracture, plate and screw fixation. Review of the literature on a case report. *Rev Cuba Reumatol.* 2019; 21(1 Supp. 1): 70.
24. Moya Rosa EJ, Marrero Peláez N. Thoracic trauma. A four-year study. *Rev Arch Méd Camagüey.* 2002; 6(6): 602-608.
25. von Garrel T, Ince A, Junge A, Schnabel M, Bahrs C. The sternal fracture: Radiographic analysis of 200 fractures with special reference to concomitant injuries. *J Trauma.* 2004; 57(4): 837-844. doi: [10.1097/01.ta.0000091112.02703.d8](https://doi.org/10.1097/01.ta.0000091112.02703.d8)
26. Şimşek S, Özmen CA, Onat S. Morbidity and mortality associated with fracture of the sternum due to blunt trauma, by fracture type and location. *Radiol Bras.* 2022; 55(3): 167-172. doi: [10.1590/0100-3984.2021.0074](https://doi.org/10.1590/0100-3984.2021.0074)
27. Ramgopal S, Shaffey SA, Conti KA. Pediatric sternal fractures from a Level 1 trauma center. *J Pediatr Surg.* 2019; 54(8): 1628-1631. doi: [10.1016/j.jpedsurg.2018.08.040](https://doi.org/10.1016/j.jpedsurg.2018.08.040)
28. Kim EY, Yang HJ, Sung YM, Hwang KH, Kim JH, Kim HS. Sternal fracture in the emergency department: Diagnostic value of multidetector CT with sagittal and coronal reconstruction images. *Eur J Radiol.* 2012; 81(5): e708-e711. doi: [10.1016/j.ejrad.2011.05.029](https://doi.org/10.1016/j.ejrad.2011.05.029)
29. Dua A, McMaster J, Desai PJ, et al. The Association between blunt cardiac injury and isolated sternal fracture. *Cardiol Res Pract.* 2014; 2014: 629687. doi: [10.1155/2014/629687](https://doi.org/10.1155/2014/629687)
30. Potaris K, Gakidis J, Mihos P, Voutsinas V, Deligeorgis A, Petsinis V. Management of sternal fractures: 239 cases. *Asian Cardiovasc Thorac Ann.* 2002; 10(2): 145-149. doi: [10.1177/021849230201000212](https://doi.org/10.1177/021849230201000212)
31. O'Connor JV, Adamski J. The diagnosis and treatment of non-cardiac thoracic trauma. *J R Army Med Corps.* 2010; 156(1): 5-14. doi: [10.1136/jramc-156-01-02](https://doi.org/10.1136/jramc-156-01-02)
32. Sarquis G, Vejez SE, Suizer A, Reche F. Fracturas traumáticas del esternón: Opciones de diagnóstico y tratamiento / Diagnostic and therapeutic options for traumatic sternal fractures. [In: Spanish]. *Rev Fac Cienc Méd (Córdoba).* 2003; 60(1): 13-18.
33. Salas VP, Bannura YF, Solovera RME, et al. Sternal fracture: Description of surgical treatment. *Rev Cir.* 2021; 73(6): 758-762.
34. Harston A, Roberts C. Fixation of sternal fractures: A systematic review. *J Trauma.* 2011; 71(6): 1875-1879. doi: [10.1097/TA.0b013e31823c46e8](https://doi.org/10.1097/TA.0b013e31823c46e8)