

Systematic Review

*Corresponding author

Laisa Liane Paineiras-Domingos, MSc
Laboratório de Vibrações Mecânicas e
Práticas Integrativas e Complementares
Departamento de Biofísica e Biometria
Instituto de Biologia Roberto Alcântara
Gomes, Universidade do Estado do Rio
de Janeiro, Av. 28 de Setembro
87, fundos, 4º andar, 20551031, Rio de
Janeiro, RJ, Brasil
Tel. 55 21 28688332
E-mail: laisanit@gmail.com

Volume 2 : Issue 1

Article Ref. #: 1000TCOJ1106

Article History

Received: January 17th, 2017

Accepted: February 27th, 2017

Published: February 27th, 2017

Citation

Sá-Caputo D, Paineiras-Domingos LL, Guedes-Aguiar EO, et al. Could hair loss be a relevant symptom associated with chikungunya?. *Trichol Cosmetol Open J.* 2017; 2(1): 5-10. doi: [10.17140/TCOJ-2-106](https://doi.org/10.17140/TCOJ-2-106)

Copyright

©2017 Paineiras-Domingos LL. This is an open access article distributed under the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Could Hair Loss be a Relevant Symptom Associated with Chikungunya?

Danúbia Sá-Caputo, PhD^{1,2}; Laisa Liane Paineiras-Domingos, MSc^{1,2}; Eliane O. Guedes-Aguiar, MSc^{2,4}; Rafaelle P. Carvalho-Lima, MSc^{2,3}; Severo de Paoli, PhD²; Danielle Soares Morel, BSc^{1,2}; Igor Bastos, BSc²; Clailson H. A. Farias, BSc²; Cláudia F. Figueiredo, BSc²; Glenda Dias, BSc²; Eric H. F. Frederico, MSc^{2,5}; Mario Bernardo-Filho, PhD²

¹Programa de Pós-Graduação em Ciências Médicas, Universidade do Estado do Rio de Janeiro, Rio de Janeiro, RJ, Brasil

²Laboratório de Vibrações Mecânicas e Práticas Integrativas e Complementares, Departamento de Biofísica e Biometria, Instituto de Biologia Roberto Alcântara Gomes, Universidade do Estado do Rio de Janeiro, Rio de Janeiro, RJ, Brasil

³Mestrado Profissional em Saúde, Medicina Laboratorial e Tecnologia Forense, Universidade do Estado do Rio de Janeiro, Rio de Janeiro, RJ, Brasil

⁴Programa de Pós Graduação em Ciências da Saúde, Universidade Federal do Rio Grande do Norte, Natal, RN, Brasil

⁵Programa de Pós Graduação em Biociências, Universidade do Estado do Rio de Janeiro, Rio de Janeiro, RJ, Brasil

ABSTRACT

Chikungunya (CHIK) is a disease caused by a mosquito-borne alphavirus (the mosquitoes *Aedes aegypti* or *Aedes albopictus*). The chikungunya virus (CHIKV) is present all over the world. Several clinical symptoms have been associated with CHIK including hair loss. The aim of this review is to present some symptoms related to CHIK, tissues and organs targeted by the CHIKV and to suggest how the infection can lead to hair loss. Searches were performed to verify the number of publications (NP) with the keywords “Chikungunya” and (i) “fever, or arthralgia, or rash, or myalgia, or fatigue, or depression, or insomnia, or pruritus, or patches, nausea, or vomiting, or headache, or erythema or hair loss” and (ii) some organs or tissues that are target to the arboviral alphaviruses associated with this disease. The total number of articles with the keyword “chikungunya” was 3176. Fever was the most cited symptoms and 50% of the publications were related to this symptom. Also, hair loss was found to be associated with CHIKV infection. Skin being the most cited organ, the total number of publications with “skin” under chikungunya summed up to only 2.77%. Although, fever is the most reported, arthralgia and rash have an important NP, and (ii) the main targets of CHIKV would be skin, kidneys and brain. A small NP is found with CHIK and hair loss. Hair loss may be idiopathic, and would be associated with a genetic predisposition, metabolic and hormonal abnormalities and aging. Also, as hair does not have a vital function in humans, it is possible to say that hair loss could not be considered in the clinical evaluation of the patient with CHIK. It is therefore possible to conclude that occurrence of hair loss is underestimated in CHIK.

KEYWORDS: Chikungunya; Hair loss; *Aedes* mosquitoes; *Aedes aegypti*; *Aedes albopictus*.

INTRODUCTION

Chikungunya (CHIK) is a disease caused by a mosquito-borne alphavirus, called the Chikungunya virus (CHIKV). It is found that CHIKV circulates in two distinct transmission cycles (enzootic transmission and urban transmission). The enzootic transmission occurs among non-human primates and perhaps other vertebrates by arboreal *Aedes* spp. mosquitoes in sub-Saharan African sylvatic foci. The urban transmission occurs in humans by *Aedes* mosquitoes (*Aedes aegypti*, *Aedes albopictus*).^{1,2} CHIKV is an enveloped, RNA positive-strand alphavirus belonging to the *Togaviridae* family.³

Figure 1 shows the relation between CHIK and the CHIKV, as well as, the forms of transmission by the *Aedes* mosquitoes.

CHIKV is an African virus that circulates enzootically in a sylvatic cycle between arboreal, canopy-dwelling mosquitoes and non-human primates.² CHIK is an endemic disease to many countries in Western, Central, Eastern and Southern Africa, Indian Ocean, West Pacific Islands and South-East Asia.¹ Since 2004, CHIK has expanded its geographic range to Indian Ocean Islands, Italy, France, and America.^{1,2}

Despite several clinical symptoms being associated with CHIK,^{2,4} little attention is paid to scalp lesions in CHIKV infection that can lead to hair loss.⁴ This could be related to the fact that hair has no vital function in humans; although, it has important psychological functions that are recognizable in most people.⁵ However, in the field of alpha viral infections, hair loss is consistent with alopecia and was reported in an experimental model with mice.⁶ These symptoms related to CHIK are associated with several organs and tissues. Based on the study by Soumahoro et al⁴ (Figure 2), certain general symptoms as well as organs and tissues which could be affected by CHIK are mentioned. Also the clinical approaches associated with hair loss have been specified (Figure 2).

Turner et al⁷ consider that hair is the ultimate personal beauty tool of self-expression. In addition, it is, in general, more malleable than skin and it is more personal than clothing. Hunt and McHale⁸ have described that alopecia is a chronic dermatological disorder in which hair fall occurs partly or completely from the areas such as the scalp or other parts of the body.

Based on the study by Hunt and McHale⁸ alopecia is considered as a chronic inflammatory disease that affects the hair follicles. It is neither life threatening nor painful; however, there can be irritation of the skin, as well as appearance related problems resulting from the loss of eyelashes and eyebrows. Alopecia can also be associated with serious psychological consequences, particularly in relation to anxiety and depression. Wattras et al⁹ reported that hair loss might also be associated with the use of medications, such as anticoagulants. Different medications have been used for the treatment of CHIKV infections.^{10,11}

Some factors associated with hair loss are found to be related to CHIK, as the CHIKV can target different human cells, such as epithelial and endothelial cells, fibroblasts and macrophages,¹² HeLa cells¹³ and muscle progenitor cells.¹⁴ Therefore, CHIKV can be associated with clinical manifestations including rashes¹⁵⁻¹⁷ and urticaria.¹⁸ CHIKV could also damage collagen and matrix metalloproteinases (MMPs)¹⁹ and alter the connective tissue metabolism.²⁰ In skeletal muscles, the atrophy and necrosis of scattered muscle fibers and the vacuolization of cells have also been shown to have potential to affect underlying mechanisms related to a disorder known as myositis syndrome.¹² Putting together these findings, it is relevant to consider that if these phenomena can lead to degenerative chronic lesions, it is possible to say that the target human epithelial and endothelial cells, fibroblasts and macrophages,¹² muscle progenitor cells,¹⁴ hypermelanosis, xerosis¹⁸ might cause a great variety of clinical manifestations including hair loss.⁴

Various authors²¹⁻²³ have used databases, such as PubMed, to evaluate the scientific findings in this topic. PubMed is a database that comprises of more than 26 million citations

Figure 1: Forms of transmission of the Chikungunya virus by the *Aedes* mosquitoes.





Figure 2: General Approaches of the Chikungunya Disease and the Hair Loss.

for biomedical literature from MEDLINE, life sciences journals, and online books. Citations may include links to full-text content from PubMed Central and publisher websites. Several studies have utilized this database for revisions and investigations involving CHIK.²¹⁻²³

The aim of this review is to present some symptoms related to CHIK and tissues and organs that are target to the CHIKV, leading to hair loss.

MATERIAL AND METHODS

Database used:

Two searches were performed through PubMed (<http://www.ncbi.nlm.nih.gov/pubmed>) database.

1. Search strategy used to verify the publications involving chikungunya and some symptoms associated with this disease caused by an arboviral alphavirus.

The searches were performed to verify the number of publications (NP) with the keywords, "chikungunya and fever", or "arthralgia", or "rash", or "myalgia", or "fatigue", or "depression", or "insomnia", or "pruritus", or "patches", "nausea", or "vomiting", or "headache", or "erythema" or "hair loss".

2. Search strategy used to verify the publications involving chikungunya and some organs or tissues that are target of the arboviral alphavirus associated with this disease.

The searches were performed to verify the NP with the keywords "chikungunya and heart", or "liver", or "kidneys", or "intestines", or "skin", or "brain", or "muscle", or "bone", or "cartilage" or "collagen".

Three reviewers independently accessed both the bibliographical databases through the *Universidade do Estado do Rio de Janeiro* on January 5th 2017.

Inclusion and Exclusion criteria to select the publications:

All the publications about chikungunya and "hair loss" that were found in the PubMed database were included in this narrative review.

RESULTS

Table 1 shows the number and the percentage of the publications related to Chikungunya, that is an endemic disease in some countries. It also shows the symptoms and percentage of each symptom recorded in the number of publications with this infection in the PubMed database. The total number of articles with the keyword "Chikungunya" is 3,176. It is observed that fever is the most cited symptom and 50% of the publications are related to this symptom. Arthralgia and rashes are other disorders associated with Chikungunya as shown in a relevant number of articles. It is also observed that hair loss is also a symptom that would be associated with CHIKV infection; however, it is poorly studied, with only one publication in the PubMed database.

Table 1: Number of the Publications Involving Chikungunya and some Symptoms and the Percentage of each Symptom Related to the Total Number of Publications with this Infection in the PubMed Database.

Symptoms	Number of publications	Percentage of the publications (%)
Fever	1611	50.72
Arthralgia	283	8.91
Rash	235	7.40
Myalgia	75	2.36
Headache	71	2.24
Fatigue	22	0.69
Nausea	22	0.69
Vomiting	21	0.66
Depression	14	0.44
Patches	8	0.25
Erythema	7	0.22
Pruritus	6	0.19
Insomnia	1	0.03
"hair loss"	1	0.03

Table 2: Number and Percentage of the Publications Involving Chikungunya and Some Targets (organs/tissues) to the Chikungunya Virus in the PubMed Database.

Organ/tissues	Number of publications	Percentage of the publications (%)
Skin	88	2.77
Kidney	68	2.14
Brain	64	2.02
Muscle	49	1.54
Bone	29	0.91
Liver	22	0.69
Heart	9	0.28
Intestine	7	0.22
Cartilage	7	0.22
Collagen	4	0.13

Table 2 shows the number of the publications related to Chikungunya and some targets (organs or tissues) with its percentage related to the total number of publications. Here, it is observed that skin is the most cited organ, but with only 2.77% of the total of publications with Chikungunya. It is also verified that the number of studies with the association between Chikungunya and kidney are high too.

DISCUSSION

Following the aim of this review and considering the number of publications in PubMed, (i) amongst the symptoms related to CHIK, fever is the most reported, whereas, arthralgia and rash have significant NP (ii) the main targets of CHIKV were found to be skin, kidney and brain. Very few publications have been found with CHIK and hair loss. These finding could be associated with the fact that the hair does not have a vital function in

humans, as reported by Ramos-e-Silva et al.⁵ In this case, it is possible to say that hair loss could not be considered in the clinical evaluation of the patient with CHIK, although, Mirmirani,²⁴ considers that the hair is one of the most defining aspects of human appearance. Moreover, hair loss may be idiopathic, and would be associated with a genetic predisposition, metabolic and hormonal abnormalities and aging.^{5,7}

The basis for this symptom following CHIKV infection is not understood. It could simply reflect a non-specific symptom triggered by the acute febrile illness, as reported by Ramos-e-Silva et al.⁵ after many systemic diseases. Although, various clinical symptoms have been associated with CHIK,^{2,4} little attention is paid to scalp lesions in CHIKV infection that can lead to the hair loss. Soumahoro et al.⁴ reported a comparative study of longstanding disabilities owing to CHIKV infection. It was showed several subjective symptoms (arthralgia, myalgia,

fatigue, depression, hair loss) reported by people infected by CHIKV.

According to Hunt and McHale,⁸ alopecia is an inflammatory disease that affects the hair follicles. This consideration could help to understand the findings presented in Table 1. The analysis of the results reported in Table 1 indicates possible symptoms that could contribute directly to hair loss, as well as rashes, pruritus, patches and erythema. Moreover, other symptoms, such as depression and insomnia could be also associated with alopecia. As the hair is not considered a vital structure in the human beings, hair loss could not be an essential question to be asked in the clinical evaluation of the individual with CHIK disease.

By analyzing the findings reported in Table 2, it is possible verify the importance of certain tissues in which, some alteration could lead to hair loss, as well as affect the skin, muscle and collagen. Authors have suggested that CHIKV can cause a wide range of clinical manifestations including rashes, hypermelanosis and xerosis¹⁶⁻¹⁸ that can further contribute to hair loss. Moreover, damaged collagen, altered connective tissue metabolism, degenerative chronic lesions by the CHIKV could also be related to alopecia.^{4,14,20} It is important to point out that medications could also sometimes promote hair loss, as reported by Yang et al²⁵ in the treatment of chronic hepatitis.

As the CHIKV infection can target different types of human cells^{12,14} and lead to many injuries in the collagen, in the matrix metalloproteinases,¹⁹ in the connective tissue metabolism,²⁰ in the skeletal muscles, degenerative chronic lesions related to inflammatory responses and the possible subsequent effect of immune activation could promote hair loss.

All the above findings, showing a relation between CHIKV infection and hair loss, are reinforced with experimental studies. Ziegler et al⁶ described that newborn mice inoculated subcutaneously with CHIKV showed signs of illness (lethargy, loss of balance and difficulty in walking, dragging of the hind limbs, and hair loss around the inoculation site on the back) 7-10 days after infection. Vanlandingham et al²⁶ reported that O'nyong-nyong virus (ONNV) and CHIKV are closely related viruses in the Semliki Forest antigenic complex (family Togaviridae, genus *Alphavirus*), whereas Walker et al²⁷ reported that mice that were inoculated with unadapted ONNV at 28-30 hours of age, developed a patchy alopecia on the 4th-7th post-inoculation day, which usually lasted 4-5 days.

The current review has certain limitations that must be considered in the interpretation of the findings. Firstly, it includes only the papers found in the PubMed database. In addition, the keywords used in the searches were related to the symptoms, tissues and organs and Chikungunya identified in Soumahoro et al⁴ Therefore, more revision with other databases and keywords would be instrumental.

CONCLUSION

In conclusion, despite the limitations, it can be stated that, fever and skin are the two highly emerging terms related to Chikungunya as observed in a large number of publications. Moreover, it was observed that the presence of various symptoms mentioned above could lead to alopecia in patients with Chikungunya. In addition, as hair does not have a vital function in humans, it is possible to conclude that hair loss cannot be considered in the clinical evaluation of the patient with CHIK. Hence, hair loss cannot be yet accounted as a symptom of Chikungunya.

ACKNOWLEDGMENTS

The authors thank the Brazilian Agencies (CNPq and FAPERJ) for the support.

CONFLICTS OF INTEREST

The authors declare no conflicts of interest.

AUTHORS' CONTRIBUTION

DSC, LLPD, EOGA, RPCL, SP, DSM, IB, CAF, CF, GD, EF and MBF participated in the elaboration and execution of the study. DSC, LLPD and EOGA accessed the bibliographical database. All the authors analyzed and discussed the data of the study. DSC, LLPD and MBF wrote the manuscript. EOGA, RPCL, DSM, IB, CAF, CF, GD and MBF reviewed the final version of the manuscript. All the authors read and approved the final version of this investigation.

REFERENCES

1. Powers AM, Logue CH. Changing patterns of chikungunya virus: Re-emergence of a zoonotic arbovirus. *J Gen Virol.* 2007; 88: 2363-2377. doi: [10.1099/vir.0.82858-0](https://doi.org/10.1099/vir.0.82858-0)
2. Coffey LL, Failloux AB, Weaver SC. Chikungunya virus-vector interactions. *Viruses.* 2014; 24: 4628-4663.
3. Porterfield JH. Antigenic characteristics and classification of the Togaviridae. In: Schlesinger R, ed. *The Togaviruses*. New York, USA: Academic Press; 1980: 13-46.
4. Soumahoro MK, Gérardin P, Boëlle PY, et al. Impact of Chikungunya virus infection on health status and quality of life: A retrospective cohort study. *PLoS One.* 2009; 4: e7800. doi: [10.1371/journal.pone.0007800](https://doi.org/10.1371/journal.pone.0007800)
5. Ramos-e-Silva M, Chaves Azevedo-e-Silva M, Carneiro SC. Hair, nail, and pigment changes in major systemic disease. *Clin Dermatol.* 2008; 26(3): 296-305. doi: [10.1016/j.clindermatol.2007.10.019](https://doi.org/10.1016/j.clindermatol.2007.10.019)
6. Ziegler SA, Lu L, da Rosa AP, Xiao SY, Tesh RB. An animal

- model for studying the pathogenesis of chikungunya virus infection. *Am J Trop Med Hyg.* 2008; 79(1): 133-139. Web site. <http://www.ajtmh.org/content/79/1/133.long>. Accessed January 16, 2017.
7. Turner GA, Bhogal RK. Hair and Aging. *Skinmed.* 2016; 14(5): 338-343.
8. Hunt N, McHale S. The psychological impact of alopecia. *BMJ.* 2005; 331: 951-953. Web site. <http://search.proquest.com/openview/b92307a5dc9d8cdb21b0bfe85268beab/1?pq-origsite=gscholar&cbl=2040978>. Accessed January 16, 2017.
9. Watras MM, Patel JP, Arya R. Traditional anticoagulants and hair loss: A role for direct oral anticoagulants? A review of the literature. *Drugs Real World Outcomes.* 2016; 3(1): 1-6. doi: [10.1007/s40801-015-0056-z](https://doi.org/10.1007/s40801-015-0056-z)
10. Gallegos KM, Drusano GL, D Argenio DZ, Brown AN. Chikungunya Virus: In Vitro Response to Combination Therapy With Ribavirin and Interferon Alfa 2a. *J Infect Dis.* 2016; 214(8): 1192-1197. doi: [10.1093/infdis/jiw358](https://doi.org/10.1093/infdis/jiw358)
11. Abdelnabi R, Neyts J, Delang L. Antiviral Strategies Against Chikungunya Virus. *Methods Mol Biol.* 2016; 1426: 243-253. doi: [10.1007/978-1-4939-3618-2_22](https://doi.org/10.1007/978-1-4939-3618-2_22)
12. Sourisseau M, Schilte C, Casartelli N, et al. Characterization of reemerging chikungunya virus. *PLoS Pathog.* 2007; 3: e89. doi: [10.1371/journal.ppat.0030089](https://doi.org/10.1371/journal.ppat.0030089)
13. Krejbich-Trotot P, Denizot M, Hoarau JJ, Jaffar-Bandjee MC, Das T, Gasque P. Chikungunya virus mobilizes the apoptotic machinery to invade host cell defenses. *FASEB J.* 2011; 25(1): 314-325. doi: [10.1096/fj.10-164178](https://doi.org/10.1096/fj.10-164178)
14. Ozden S, Huerre M, Riviere JP, et al. Human muscle satellite cells as targets of Chikungunya virus infection. *PLoS ONE.* 2007; 2: e527. doi: [10.1371/journal.pone.0000527](https://doi.org/10.1371/journal.pone.0000527)
15. Cunha BA, Leonichev VB, Raza M. Chikungunya fever presenting with protracted severe pruritus. *IDCases.* 2016; 6: 29-30. doi: [10.1016/j.idcr.2016.09.003](https://doi.org/10.1016/j.idcr.2016.09.003)
16. Borgherini G, Poubeau P, Jossaume A, et al. Persistent arthralgia associated with chikungunya virus: A study of 88 adult patients on reunion island. *Clin Infect Dis.* 2008; 47: 469-475. doi: [10.1086/590003](https://doi.org/10.1086/590003)
17. Borgherini G, Poubeau P, Staikowsky F, et al. Outbreak of chikungunya on Reunion Island: early clinical and laboratory features in 157 adult patients. *Clin Infect Dis.* 2007; 44(11): 1401-1407. doi: [10.1086/517537](https://doi.org/10.1086/517537)
18. Bandyopadhyay D, Ghosh SK. Mucocutaneous features of Chikungunya fever: A study from an outbreak in West Bengal, India. *Int J Dermatol.* 2008; 47(11): 1148-1152. doi: [10.1111/j.1365-4632.2008.03817.x](https://doi.org/10.1111/j.1365-4632.2008.03817.x)
19. Kantor AM, Dong S, Held NL, et al. Identification and initial characterization of matrix metalloproteinases in the yellow fever mosquito, *Aedes aegypti*. *Insect Mol Biol.* 2017; 26(1): 113-126. doi: [10.1111/imb.12275](https://doi.org/10.1111/imb.12275)
20. Lokireddy S, Vemula S, Vadde R. Connective tissue metabolism in chikungunya patients. *Virology.* 2008; 5: 31. doi: [10.1186/1743-422X-5-31](https://doi.org/10.1186/1743-422X-5-31)
21. Rodríguez-Morales AJ, Cardona-Ospina JA, Fernanda Urbano-Garzón S, Sebastian Hurtado-Zapata J. Prevalence of Post-Chikungunya Infection Chronic Inflammatory Arthritis: A Systematic Review and Meta-Analysis. *Arthritis Care Res (Hoboken).* 2016; 68: 1849-1858. doi: [10.1002/acr.22900](https://doi.org/10.1002/acr.22900)
22. Furuya-Kanamori L, Liang S, Milinovich G, et al. Co-distribution and co-infection of chikungunya and dengue viruses. *BMC Infect Dis.* 2016; 16: 84. doi: [10.1186/s12879-016-1417-2](https://doi.org/10.1186/s12879-016-1417-2)
23. Thiberville SD, Moyen N, Dupuis-Maguiraga L, et al. Chikungunya fever: Epidemiology, clinical syndrome, pathogenesis and therapy. *Antiviral Res.* 2013; 99(3): 345-370. doi: [10.1016/j.antiviral.2013.06.009](https://doi.org/10.1016/j.antiviral.2013.06.009)
24. Mirmirani P. Managing hair loss in midlife women. *Maturitas.* 2013; 74(2): 119-122. doi: [10.1016/j.maturitas.2012.10.020](https://doi.org/10.1016/j.maturitas.2012.10.020)
25. Yang SY, Lee HW, Lee YJ, Park SJ, Yoo KY, Kim HJ. Highly effective peginterferon α -2a plus ribavirin combination therapy for chronic hepatitis C in hemophilia in Korea. *Clin Mol Hepatol.* 2015; 21(2): 125-130. doi: [10.3350/cmh.2015.21.2.125](https://doi.org/10.3350/cmh.2015.21.2.125)
26. Vanlandingham DL, Hong C, Klingler K, et al. Differential infectivities of o'nyong-nyong and chikungunya virus isolates in *Anopheles gambiae* and *Aedes aegypti* mosquitoes. *Am J Trop Med Hyg.* 2005; 72(5): 616-621. Web site. <http://www.ajtmh.org/content/72/5/616.long>. Accessed January 16, 2017.
27. Walker GM, Woodall JP, Haddow AJ, Williams MC. O'nyong-nyong fever: An epidemic virus disease in East Africa. VI. Alopecia in mice experimentally infected with o'nyong-nyong virus. *Trans R Soc Trop Med Hyg.* 1962; 56(6): 496-503. doi: [10.1016/0035-9203\(62\)90073-1](https://doi.org/10.1016/0035-9203(62)90073-1)